

# Alligator Harbor Aquatic Preserve

## SEACAR Habitat Analyses

Last compiled on 08 January, 2025

### Contents

<b>Funding &amp; Acknowledgements</b>	<b>2</b>
<b>Threshold Filtering</b>	<b>2</b>
<b>Value Qualifiers</b>	<b>3</b>
<b>Water Column</b>	<b>5</b>
<b>Seasonal Kendall-Tau Analysis</b>	<b>5</b>
<b>Water Quality - Discrete</b>	<b>5</b>
Chlorophyll a, Uncorrected for Pheophytin - Discrete Water Quality . . . . .	6
Colored Dissolved Organic Matter - Discrete Water Quality . . . . .	7
Dissolved Oxygen - Discrete Water Quality . . . . .	9
pH - Discrete Water Quality . . . . .	12
Salinity - Discrete Water Quality . . . . .	14
Secchi Depth - Discrete Water Quality . . . . .	16
Total Nitrogen - Discrete Water Quality . . . . .	18
Total Phosphorus - Discrete Water Quality . . . . .	20
Turbidity - Discrete Water Quality . . . . .	22
Water Temperature - Discrete Water Quality . . . . .	24
<b>Water Quality - Continuous</b>	<b>28</b>
Dissolved Oxygen - All Stations Combined . . . . .	30
Dissolved Oxygen Saturation - All Stations Combined . . . . .	31
pH - All Stations Combined . . . . .	32
Salinity - All Stations Combined . . . . .	33
Turbidity - All Stations Combined . . . . .	34
Water Temperature - All Stations Combined . . . . .	35
<b>Submerged Aquatic Vegetation</b>	<b>36</b>
Parameters . . . . .	36
Species . . . . .	36
Notes . . . . .	36
<b>References</b>	<b>42</b>

## Funding & Acknowledgements

The data used in this analysis is from the Export Standardized Tables in the SEACAR Data Discovery Interface (DDI). Documents and information available through the SEACAR DDI are owned by the data provider(s) and users are expected to provide appropriate credit following accepted citation formats. Users are encouraged to access data to maximize utilization of gained knowledge, reducing redundant research and facilitating partnerships and scientific innovation.

With respect to documents and information available from SEACAR DDI, neither the State of Florida nor the Florida Department of Environmental Protection makes any warranty, expressed or implied, including the warranties of merchantability and fitness for a particular purpose arising out of the use or inability to use the data, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

This report was funded in part, through a grant agreement from the Florida Department of Environmental Protection, Florida Coastal Management Program, by a grant provided by the Office for Coastal Management under the Coastal Zone Management Act of 1972, as amended, National Oceanic and Atmospheric Administration. The views, statements, findings, conclusions and recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of the State of Florida, NOAA or any of their sub agencies.

Published: 2025-01-08



## Threshold Filtering

Threshold filters, following the guidance of Florida Department of Environmental Protection’s (*FDEP*) Division of Environmental Assessment and Restoration (*DEAR*) are used to exclude specific results values from the SEACAR Analysis. Based on the threshold filters, Quality Assurance / Quality Control (*QAQC*) Flags are inserted into the *SEACAR\_QAQCFlagCode* and *SEACAR\_QAQC\_Description* columns of the export data. The *Include* column indicates whether the *QAQC* Flag will also indicate that data are excluded from analysis. No data are excluded from the data export, but the analysis scripts can use the *Include* column to exclude data (1 to include, 0 to exclude).

Table 1: Continuous Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	-0.000001	50
Dissolved Oxygen Saturation	%	-0.000001	500
Salinity	ppt	-0.000001	70
Turbidity	NTU	-0.000001	4000
Water Temperature	Degrees C	-5.000000	45
pH	None	2.000000	14

Table 2: Discrete Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Ammonia, Un-ionized (NH3)	mg/L	-	-
Ammonium, Filtered (NH4)	mg/L	-	-
Chlorophyll a, Corrected for Pheophytin	ug/L	-	-
Chlorophyll a, Uncorrected for Pheophytin	ug/L	-	-
Colored Dissolved Organic Matter	PCU	-	-

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	-0.000001	25
Dissolved Oxygen Saturation	%	-0.000001	310
Fluorescent Dissolved Organic Matter	QSE	-	-
Light Extinction Coefficient	m <sup>-1</sup>	-	-
NO <sub>2</sub> + <sub>3</sub> , Filtered	mg/L	-	-
Nitrate (NO <sub>3</sub> )	mg/L	-	-
Nitrite (NO <sub>2</sub> )	mg/L	-	-
Nitrogen, organic	mg/L	-	-
Phosphate, Filtered (PO <sub>4</sub> )	mg/L	-	-
Salinity	ppt	-0.000001	70
Secchi Depth	m	0.000001	50
Specific Conductivity	mS/cm	0.005000	100
Total Kjeldahl Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Phosphorus	mg/L	-	-
Total Suspended Solids	mg/L	-	-
Turbidity	NTU	-	-
Water Temperature	Degrees C	3.000000	40
pH	None	2.000000	13

Table 3: Quality Assurance Flags inserted based on threshold checks listed in Table 1 and 2

<i>SEACAR QAQC Description</i>	<i>Include</i>	<i>SEACAR QAQCFlagCode</i>
Exceeds maximum threshold	0	2Q
Below minimum threshold	0	4Q
Within threshold tolerance	1	6Q
No defined thresholds for this parameter	1	7Q

## Value Qualifiers

Value qualifier codes included within the data are used to exclude certain results from the analysis. The data are retained in the data export files, but the analysis uses the *Include* column to filter the results.

### STORET and WIN value qualifier codes

Value qualifier codes from *STORET* and *WIN* data are examined with the database and used to populate the *Include* column in data exports.

Table 4: Value Qualifier codes excluded from analysis

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>MDL</i>	<i>Description</i>
STORET-WIN	H	0	0	Value based on field kit determination; results may not be accurate
STORET-WIN	J	0	0	Estimated value
STORET-WIN	V	0	0	Analyte was detected at or above method detection limit
STORET-WIN	Y	0	0	Lab analysis from an improperly preserved sample; data may be inaccurate

### Discrete Water Quality Value Qualifiers

The following value qualifiers are highlighted in the Discrete Water Quality section of this report. An exception is made for **Program 476 - Charlotte Harbor Estuaries Volunteer Water Quality Monitoring Network** and data flagged with Value Qualifier **H** are included for this program only.

**H** - Value based on field kit determination; results may not be accurate. This code shall be used if a field screening test (e.g., field gas chromatograph data, immunoassay, or vendor-supplied field kit) was used to generate the value and the field kit or method has not been recognized by the Department as equivalent to laboratory methods.

**I** - The reported value is greater than or equal to the laboratory method detection limit but less than the laboratory practical quantitation limit.

**Q** - Sample held beyond the accepted holding time. This code shall be used if the value is derived from a sample that was prepared or analyzed after the approved holding time restrictions for sample preparation or analysis.

**S** - Secchi disk visible to bottom of waterbody. The value reported is the depth of the waterbody at the location of the Secchi disk measurement.

**U** - Indicates that the compound was analyzed for but not detected. This symbol shall be used to indicate that the specified component was not detected. The value associated with the qualifier shall be the laboratory method detection limit. Unless requested by the client, less than the method detection limit values shall not be reported

**Systemwide Monitoring Program (SWMP) value qualifier codes**

Value qualifier codes from the *SWMP* continuous program are examined with the database and used to populate the *Include* column in data exports. *SWMP* Qualifier Codes are indicated by *QualifierSource=SWMP*.

Table 5: SWMP Value Qualifier codes

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>Description</i>
SWMP	-1	Yes	Optional parameter not collected
SWMP	-2	No	Missing data
SWMP	-3	No	Data rejected due to QA/QC
SWMP	-4	No	Outside low sensor range
SWMP	-5	No	Outside high sensor range
SWMP	0	Yes	Passed initial QA/QC checks
SWMP	1	No	Suspect data
SWMP	2	Yes	Reserved for future use
SWMP	3	Yes	Calculated data: non-vented depth/level sensor correction for changes in barometric pressure
SWMP	4	Yes	Historical: Pre-auto QA/QC
SWMP	5	Yes	Corrected data

## Water Column

The water column habitat extends from the water's surface to the bottom sediments, and it's where fish, dolphins, crabs and people swim! So much life makes its home in the water column that the health of marine and coastal ecosystems, as well as human economies, depend on the condition of this vulnerable habitat. Local patterns of rainfall, temperature, winds and currents can rapidly change the condition of the water column, while global influences such as [El Niño/La Niña](#), large-scale fluctuation in sea temperatures and climate change can have long-term effects. Inputs from the prosperity of our day-to-day lives including farming, mining and forestry, and emissions from power generation, automobiles and water treatment can also alter the health of the water column. Acting alone or together, each input can have complex and lasting effects on habitats and ecosystems.

SEACAR evaluates water column health with several essential parameters. These include nutrient surveys of nitrogen and phosphorus, and water quality assessments of salinity, dissolved oxygen, pH, and water temperature. Water clarity is evaluated with Secchi depth, turbidity, levels of chlorophyll a, total suspended solids, and colored dissolved organic matter. Additionally, the richness of nekton is indicated by the abundance of free-swimming fishes and macroinvertebrates like crabs and shrimps.

## Seasonal Kendall-Tau Analysis

Indicators must have a minimum of five to ten years, depending on the habitat, of data within the geographic range of the analysis to be included in the analysis. Ten years of data are required for discrete parameters, and five years of data are required for continuous parameters. If there are insufficient years of data, the number of years of data available will be noted and labeled as “insufficient data to conduct analysis”. Further, for the preferred Seasonal Kendall-Tau test, there must be data from at least two months in common across at least two consecutive years within the RCP managed area being analyzed. Values that pass both of these tests will be included in the analysis and be labeled as *Use\_In\_Analysis* = **TRUE**. Any that fail either test will be excluded from the analyses and labeled as *Use\_In\_Analysis* = **FALSE**. The points for all Water Column plots displayed in this section are monthly averages. Trend significance will be denoted as “Significant Trend” (when  $p < 0.05$ ), or “Non-significant Trend” (when  $p \geq 0.05$ ). Any parameters with insufficient data to perform Seasonal Kendall-Tau test will have their monthly averages plotted without a corresponding trend line.

## Water Quality - Discrete

The following files were used in the discrete analysis:

- *Combined\_WQ\_WC\_NUT\_Chlorophyll\_a\_corrected\_for\_pheophytin-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Chlorophyll\_a\_uncorrected\_for\_pheophytin-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Colored\_dissolved\_organic\_matter\_CDOM-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Dissolved\_Oxygen-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Dissolved\_Oxygen\_Saturation-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_pH-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Salinity-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Secchi\_Depth-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Total\_Nitrogen-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Total\_Phosphorus-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Total\_Suspended\_Solids\_TSS-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Turbidity-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Water\_Temperature-2024-Dec-08.txt*

# Chlorophyll a, Uncorrected for Pheophytin - Discrete Water Quality

## Seasonal Kendall-Tau Trend Analysis




Figure 1: Seasonal Kendall-Tau Results for Chlorophyll a, Uncorrected for Pheophytin - Discrete

Table 6: Seasonal Kendall-Tau Trend Analysis for Chlorophyll a, Uncorrected for Pheophytin

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	1213	18	6	TRUE	0.2862	0	0.1488	3.3869	8.8982	0.6313	1




Figure 2: Map showing location of Discrete sampling sites for Chlorophyll a, Uncorrected for Pheophytin. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 7: Programs contributing data for Chlorophyll a, Uncorrected for Pheophytin

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
514	1062	2001	2024
103	132	2020	2021
5002	19	2019	2022

**Program names:**

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>

514 - Florida LAKEWATCH Program<sup>2</sup>

5002 - Florida STORET / WIN<sup>3</sup>

**Colored Dissolved Organic Matter - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**

Colored Dissolved Organic Matter - Discrete, All Depths  
Alligator Harbor Aquatic Preserve




Figure 3: Seasonal Kendall-Tau Results for Colored Dissolved Organic Matter - Discrete

Table 8: Seasonal Kendall-Tau Trend Analysis for Colored Dissolved Organic Matter

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	409	18	8	TRUE	0.2102	0.0598	0.1772	6.5939	7.1826	0.2074	0






Figure 4: Map showing location of Discrete sampling sites for Colored Dissolved Organic Matter. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 9: Programs contributing data for Colored Dissolved Organic Matter

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
514	409	2001	2024

**Program names:**

514 - Florida LAKEWATCH Program<sup>2</sup>

**Dissolved Oxygen - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 5: Seasonal Kendall-Tau Results for Dissolved Oxygen - Discrete

Table 10: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	8140	27	7.2	TRUE	-0.1664	0.0001	-0.0325	7.8071	7.6468	0.7446	-1




Figure 6: Map showing location of Discrete sampling sites for Dissolved Oxygen. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 11: Programs contributing data for Dissolved Oxygen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	5829	1998	2022
69	1320	1998	2022
469	630	2016	2024
557	191	2005	2023
95	156	2004	2018
103	15	2021	2021

**Program names:**

- 69 - Fisheries-Independent Monitoring (FIM) Program<sup>4</sup>
- 95 - Harmful Algal Bloom Marine Observation Network<sup>5</sup>
- 103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>
- 469 - Central Panhandle Aquatic Preserve WQ Monitoring<sup>6</sup>
- 557 - Central Panhandle Aquatic Preserves Seagrass Monitoring<sup>7</sup>
- 5002 - Florida STORET / WIN<sup>3</sup>

# pH - Discrete Water Quality

## Seasonal Kendall-Tau Trend Analysis




Figure 7: Seasonal Kendall-Tau Results for pH - Discrete

Table 12: Seasonal Kendall-Tau Trend Analysis for pH

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	5266	27	8.1	TRUE	-0.1855	0	-0.0075	8.1378	12.4755	0.329	-1




Figure 8: Map showing location of Discrete sampling sites for pH. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 13: Programs contributing data for pH

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	2810	1998	2022
69	1310	1998	2022
469	630	2016	2024
558	180	2008	2014
557	170	2005	2023
95	152	2008	2018
103	15	2021	2021

**Program names:**

69 - Fisheries-Independent Monitoring (FIM) Program<sup>4</sup>

95 - Harmful Algal Bloom Marine Observation Network<sup>5</sup>

103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>

469 - Central Panhandle Aquatic Preserve WQ Monitoring<sup>6</sup>

557 - Central Panhandle Aquatic Preserves Seagrass Monitoring<sup>7</sup>

## Salinity - Discrete Water Quality

### Seasonal Kendall-Tau Trend Analysis




Figure 9: Seasonal Kendall-Tau Results for Salinity - Discrete

Table 14: Seasonal Kendall-Tau Trend Analysis for Salinity

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	9377	28	31.2	TRUE	-0.1057	0.011	-0.0428	31.8939	3.837	0.9744	-1




Figure 10: Map showing location of Discrete sampling sites for Salinity. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 15: Programs contributing data for Salinity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	6810	1998	2022
69	1318	1998	2022
469	630	2016	2024
558	258	2008	2014
557	191	2005	2023
95	171	1996	2018

**Program names:**

- 69 - Fisheries-Independent Monitoring (FIM) Program<sup>4</sup>
- 95 - Harmful Algal Bloom Marine Observation Network<sup>5</sup>
- 469 - Central Panhandle Aquatic Preserve WQ Monitoring<sup>6</sup>
- 557 - Central Panhandle Aquatic Preserves Seagrass Monitoring<sup>7</sup>
- 558 - Franklin County Coastal Waters Seagrass Monitoring<sup>8</sup>
- 5002 - Florida STORET / WIN<sup>3</sup>

# Secchi Depth - Discrete Water Quality

## Seasonal Kendall-Tau Trend Analysis




Figure 11: Seasonal Kendall-Tau Results for Secchi Depth - Discrete

Table 16: Seasonal Kendall-Tau Trend Analysis for Secchi Depth

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
Surface	2857	27	1.1583	TRUE	-0.0009	1	0	1.1291	12.2104	0.348	0






Figure 12: Map showing location of Discrete sampling sites for Secchi Depth. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 17: Programs contributing data for Secchi Depth

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
69	1303	1998	2022
514	1084	2001	2024
558	331	2008	2017
557	105	2005	2023
5002	24	2019	2022
103	10	2021	2021

**Program names:**

- 69 - Fisheries-Independent Monitoring (FIM) Program<sup>4</sup>
- 103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>
- 514 - Florida LAKEWATCH Program<sup>2</sup>
- 557 - Central Panhandle Aquatic Preserves Seagrass Monitoring<sup>7</sup>
- 558 - Franklin County Coastal Waters Seagrass Monitoring<sup>8</sup>
- 5002 - Florida STORET / WIN<sup>3</sup>

## Total Nitrogen - Discrete Water Quality

### Total Nitrogen Calculation:

The logic for calculated Total Nitrogen was provided by Kevin O'Donnell and colleagues at FDEP (with the help of Jay Silvanima, Watershed Monitoring Section). The following logic is used, in this order, based on the availability of specific nitrogen components.

- 1)  $TN = TKN + NO3O2$ ;
- 2)  $TN = TKN + NO3 + NO2$ ;
- 3)  $TN = ORGN + NH4 + NO3O2$ ;
- 4)  $TN = ORGN + NH4 + NO2 + NO3$ ;
- 5)  $TN = TKN + NO3$ ;
- 6)  $TN = ORGN + NH4 + NO3$ ;

### Additional Information:

- Rules for use of sample fraction:
  - Florida Department of Environmental Protection (FDEP) report that if both “Total” and “Dissolved” components are reported, only “Total” is used. If the total is not reported, then the dissolved components are used as a best available replacement.
  - Total nitrogen calculations are done using nitrogen components with the same sample fraction, nitrogen components with mixed total/dissolved sample fractions are not used. In other words, total nitrogen can be calculated when TKN and NO3O2 are both total sample fractions, or when both are dissolved sample fractions. *Future calculations of total nitrogen values may be based on components with mixed sample fractions.*
- Values inserted into data:
  - ParameterName = “Total Nitrogen”
  - SEACAR\_QAQCFlagCode = “1Q”
  - SEACAR\_QAQC\_Description = “SEACAR Calculated”

### Seasonal Kendall-Tau Trend Analysis




Figure 13: Seasonal Kendall-Tau Results for Total Nitrogen - Discrete

Table 18: Seasonal Kendall-Tau Trend Analysis for Total Nitrogen

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	1585	19	0.19	TRUE	0.2813	0	0.0077	0.0818	3.4785	0.9828	1




Figure 14: Map showing location of Discrete sampling sites for Total Nitrogen. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 19: Programs contributing data for Total Nitrogen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
514	1135	2001	2024
5002	465	2001	2022

**Program names:**

514 - Florida LAKEWATCH Program<sup>2</sup>

5002 - Florida STORET / WIN<sup>3</sup>

**Total Phosphorus - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 15: Seasonal Kendall-Tau Results for Total Phosphorus - Discrete

Table 20: Seasonal Kendall-Tau Trend Analysis for Total Phosphorus

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	1149	19	0.0001	TRUE	0.5677	0	0.0012	-0.0213	3.1643	0.9884	2




Figure 16: Map showing location of Discrete sampling sites for Total Phosphorus. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 21: Programs contributing data for Total Phosphorus

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
514	1107	2001	2024
103	61	2020	2021
5002	24	2012	2022

**Program names:**

103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>

514 - Florida LAKEWATCH Program<sup>2</sup>

5002 - Florida STORET / WIN<sup>3</sup>

**Turbidity - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 17: Seasonal Kendall-Tau Results for Turbidity - Discrete

Table 22: Seasonal Kendall-Tau Trend Analysis for Turbidity

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	3543	18	5.3	TRUE	-0.2432	0.0012	-0.1119	6.5131	4.8127	0.9399	-1




Figure 18: Map showing location of Discrete sampling sites for Turbidity. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 23: Programs contributing data for Turbidity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	3543	1998	2022
469	259	2021	2024
557	39	2022	2023
103	5	2021	2021

**Program names:**

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>

469 - Central Panhandle Aquatic Preserve WQ Monitoring<sup>6</sup>

557 - Central Panhandle Aquatic Preserves Seagrass Monitoring<sup>7</sup>

5002 - Florida STORET / WIN<sup>3</sup>

**Water Temperature - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**






Figure 19: Seasonal Kendall-Tau Results for Water Temperature - Discrete

Table 24: Seasonal Kendall-Tau Trend Analysis for Water Temperature

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	9633	28	25.8	TRUE	-0.0467	0.2321	-0.0148	22.9438	17.2667	0.1002	0




Figure 20: Map showing location of Discrete sampling sites for Water Temperature. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 25: Programs contributing data for Water Temperature

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	7037	1998	2022
69	1324	1998	2022
469	630	2016	2024
558	271	2008	2017
557	191	2005	2023
95	166	1996	2018
103	15	2021	2021

**Program names:**

- 69 - Fisheries-Independent Monitoring (FIM) Program<sup>4</sup>
- 95 - Harmful Algal Bloom Marine Observation Network<sup>5</sup>
- 103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX<sup>1</sup>
- 469 - Central Panhandle Aquatic Preserve WQ Monitoring<sup>6</sup>
- 557 - Central Panhandle Aquatic Preserves Seagrass Monitoring<sup>7</sup>

558 - Franklin County Coastal Waters Seagrass Monitoring<sup>8</sup>  
5002 - Florida STORET / WIN<sup>3</sup>

## Water Quality - Continuous

The following files were used in the continuous analysis:

- *Combined\_WQ\_WC\_NUT\_cont\_Dissolved\_Oxygen\_NW-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_cont\_Dissolved\_Oxygen\_Saturation\_NW-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_cont\_pH\_NW-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_cont\_Salinity\_NW-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_cont\_Turbidity\_NW-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_cont\_Water\_Temperature\_NW-2024-Dec-08.txt*

### Continuous monitoring locations in Alligator Harbor Aquatic Preserve

Table 26: Central Panhandle Aquatic Preserves Continuous Water Quality Monitoring (468)

<i>ProgramLocationID</i>	<i>Years of Data</i>	<i>Use in Analysis</i>	<i>Parameters</i>
CPAH	2	FALSE	DO , DOS , pH , Sal , Turb , TempW
CPAH2	6	TRUE	DO , DOS , pH , Sal , Turb , TempW
CPFS	5	FALSE	DO , DOS , pH , Sal , Turb , TempW




Figure 21: Map showing Continuous Water Quality Monitoring sampling locations within the boundaries of Alligator Harbor Aquatic Preserve. Sites marked as *Use In Analysis* are featured in this report.

# Dissolved Oxygen - All Stations Combined




Figure 22: Figure for Dissolved Oxygen - Continuous - All stations combined

Table 27: Seasonal Kendall-Tau Results for All Stations - Dissolved Oxygen

Station	N_Data	N_Years	Period of Record	Median	tau	SennIntercept	SennSlope	p
CPAH2	89535	5	2019 - 2024	7.0	-0.17	6.49	-0.07	0.3424
CPFS	39993	4	2019 - 2023	6.4	-	-	-	-
CPAH	11557	2	2007 - 2008	8.0	-	-	-	-

## Dissolved Oxygen Saturation - All Stations Combined




Figure 23: Figure for Dissolved Oxygen Saturation - Continuous - All stations combined

Table 28: Seasonal Kendall-Tau Results for All Stations - Dissolved Oxygen Saturation

Station	N_Data	N_Years	Period of Record	Median	tau	SennIntercept	SennSlope	p
CPAH2	89535	5	2019 - 2024	94.4	-0.33	97.16	-0.94	0.0327
CPFS	39993	4	2019 - 2023	90.4	-	-	-	-
CPAH	11557	2	2007 - 2008	105.6	-	-	-	-

# pH - All Stations Combined




Figure 24: Figure for pH - Continuous - All stations combined

Table 29: Seasonal Kendall-Tau Results for All Stations - pH

Station	N_Data	N_Years	Period of Record	Median	tau	SennIntercept	SennSlope	p
CPAH2	89372	5	2019 - 2024	8.0	-0.4	8.07	-0.04	0.0241
CPAH	13034	2	2007 - 2008	7.9	-	-	-	-
CPFS	43191	4	2019 - 2023	8.1	-	-	-	-



## Salinity - All Stations Combined




Figure 25: Figure for Salinity - Continuous - All stations combined

Table 30: Seasonal Kendall-Tau Results for All Stations - Salinity

Station	N_Data	N_Years	Period of Record	Median	tau	SennIntercept	SennSlope	p
CPAH2	85969	5	2019 - 2024	30.7	-0.23	31.6	-0.13	0.2353
CPFS	38401	4	2019 - 2023	30.7	-	-	-	-
CPAH	13034	2	2007 - 2008	34.0	-	-	-	-

## Turbidity - All Stations Combined




Figure 26: Figure for Turbidity - Continuous - All stations combined

Table 31: Seasonal Kendall-Tau Results for All Stations - Turbidity

Station	N_Data	N_Years	Period of Record	Median	tau	SennIntercept	SennSlope	p
CPAH2	97620	5	2019 - 2024	5	0.22	4	0.88	0.1770
CPFS	37491	4	2019 - 2023	4	-	-	-	-
CPAH	12558	2	2007 - 2008	10	-	-	-	-

# Water Temperature - All Stations Combined




Figure 27: Figure for Water Temperature - Continuous - All stations combined

Table 32: Seasonal Kendall-Tau Results for All Stations - Water Temperature

Station	N_Data	N_Years	Period of Record	Median	tau	SennIntercept	SennSlope	p
CPFS	53708	4	2019 - 2023	24.2	-	-	-	-
CPAH2	102382	5	2019 - 2024	22.6	0.14	23.93	0.14	0.6471
CPAH	13034	2	2007 - 2008	20.9	-	-	-	-

# Submerged Aquatic Vegetation

The data file used is: **All\_SAV\_Parameters-2024-Dec-08.txt**

**Submerged aquatic vegetation (SAV)** refers to plants and plant-like macroalgae species that live entirely underwater. The two primary categories of SAV inhabiting Florida estuaries are *benthic macroalgae* and *seagrasses*. They often grow together in dense beds or meadows that carpet the seafloor. *Macroalgae* include multicellular species of green, red and brown algae that often live attached to the substrate by a holdfast. They tend to grow quickly and can tolerate relatively high nutrient levels, making them a threat to seagrasses and other benthic habitats in areas with poor water quality. In contrast, *seagrasses* are grass-like, vascular, flowering plants that are attached to the seafloor by extensive root systems. *Seagrasses* occur throughout the coastal areas of Florida, including protected bays and lagoons as well as deeper offshore waters on the continental shelf. *Seagrasses* have taken advantage of the broad, shallow shelf and clear water to produce two of the most extensive seagrass beds anywhere in continental North America.

## Parameters

**Percent Cover** measures the fraction of an area of seafloor that is covered by SAV, usually estimated by evaluating multiple small areas of seafloor. Percent cover is often estimated for total SAV, individual types of vegetation (seagrass, attached algae, drift algae) and individual species.

**Frequency of Occurrence** was calculated as the number of times a taxon was observed in a year divided by the number of sampling events, multiplied by 100. Analysis is conducted at the quadrat level and is inclusive of all quadrats (i.e., quadrats evaluated using Braun-Blanquet, modified Braun-Blanquet, and percent cover.”

## Species

**Turtle grass** (*Thalassia testudinum*) is the largest of the Florida seagrasses, with longer, thicker blades and deeper root structures than any of the other seagrasses. It is considered a climax seagrass species.

**Shoal grass** (*Halodule wrightii*) is an early colonizer of vegetated areas and usually grows in water too shallow for other species except *widgeon grass*. It can often tolerate larger salinity ranges than other seagrass species. *Shoal grass* is characterized by thin, flat blades, that are narrower than *turtle grass* blades.

**Manatee grass** (*Syringodium filiforme*) is easily recognizable because its leaves are thin and cylindrical instead of the flat, ribbon-like form shared by many other seagrass species. The leaves can grow up to half a meter in length. *Manatee grass* is usually found in mixed seagrass beds or small, dense monospecific patches.

**Widgeon grass** (*Ruppia maritima*) grows in both fresh and salt water and is widely distributed throughout Florida’s estuaries in less saline areas, particularly in inlets along the east coast. This species resembles *shoal grass* in certain environments but can be identified by the pointed tips of its leaves.

Three species of *Halophila spp.* are found in Florida - **Star grass** (*Halophila engelmannii*), **Paddle grass** (*Halophila decipiens*), and **Johnson’s seagrass** (*Halophila johnsonii*). These are smaller, more fragile seagrasses than other Florida species and are considered ephemeral. They grow along a single long rhizome, with short blades. These species are not well-studied, although surveys are underway to define their ecological roles.

## Notes

*Star grass*, *Paddle grass*, and *Johnson’s seagrass* will be grouped together and listed as **Halophila spp.** in the following managed areas. This is because several surveys did not specify to the species level:

- Banana River Aquatic Preserve
- Indian River-Malabar to Vero Beach Aquatic Preserve
- Indian River-Vero Beach to Ft. Pierce Aquatic Preserve
- Jensen Beach to Jupiter Inlet Aquatic Preserve
- Loxahatchee River-Lake Worth Creek Aquatic Preserve
- Mosquito Lagoon Aquatic Preserve

- Biscayne Bay Aquatic Preserve
- Florida Keys National Marine Sanctuary




Figure 28: Maps showing the temporal scope of SAV sampling sites within the boundaries of *Alligator Harbor Aquatic Preserve* by Program name.

**Sampling locations by Program:**




Figure 29: Map showing SAV sampling sites within the boundaries of *Alligator Harbor Aquatic Preserve*. The point size reflects the number of samples at a given sampling site.

Table 33: Franklin County Coastal Waters Seagrass Monitoring - Program 558

<i>N-Data</i>	<i>YearMin</i>	<i>YearMax</i>	<i>method</i>	<i>Sample Locations</i>
2984	2009	2017	Percent Cover	38

Table 34: Central Panhandle Aquatic Preserves Seagrass Monitoring - Program 557

<i>N-Data</i>	<i>YearMin</i>	<i>YearMax</i>	<i>method</i>	<i>Sample Locations</i>
531	2008	2023	Braun Blanquet	16




Figure 30: Trends in median percent cover for various seagrass species in Alligator Harbor Aquatic Preserve




Figure 31: Trends in median percent cover for various seagrass species in Alligator Harbor Aquatic Preserve - simplified

Table 35: Percent Cover Trend Analysis for Alligator Harbor Aquatic Preserve

<i>CommonName</i>	<i>Trend Significance (0.05)</i>	<i>Period of Record</i>	<i>LME-Intercept</i>	<i>LME-Slope</i>	<i>p</i>
Drift algae	Significantly decreasing trend	2009 - 2023	79.40406	-2.3563220	0.0273433
Shoal grass	No significant trend	2008 - 2023	29.78762	-0.4993661	0.4716922
Star grass	Insufficient data to calculate trend	-	-	-	-
No grass in quadrat	Model did not fit the available data	2009 - 2023	-	-	-
Widgeon grass	Insufficient data to calculate trend	-	-	-	-
Manatee grass	Significantly decreasing trend	2008 - 2023	47.85836	-1.2281451	0.0253568
Turtle grass	No significant trend	2008 - 2023	30.96903	-0.4530891	0.3778578



Frequency of occurrence  
Alligator Harbor Aquatic Preserve




Figure 32: Frequency of occurrence for various seagrass species in Alligator Harbor Aquatic Preserve

## References

1. U.S. Environmental Protection Agency (EPA). [EPA STORage and RETrieval Data Warehouse \(STORET\)/WQX](#). (2023).
2. University of Florida (UF); Institute of Food and Agricultural Sciences. [Florida LAKEWATCH Program](#). (2024).
3. Florida Department of Environmental Protection (DEP). [Florida STORET / WIN](#). (2024).
4. Florida Fish and Wildlife Conservation Commission (FWC). [Fisheries-Independent Monitoring \(FIM\) Program](#). (2022).
5. Florida Fish and Wildlife Conservation Commission (FWC); Florida Fish and Wildlife Research Institute (FWRI). [Harmful Algal Bloom Marine Observation Network](#). (2018).
6. Florida Department of Environmental Protection (DEP); Office of Resilience and Coastal Protection (RCP); Central Panhandle Aquatic Preserves. [Central Panhandle Aquatic Preserve WQ Monitoring](#) . (2024).
7. Florida Department of Environmental Protection (DEP); Office of Resilience and Coastal Protection (RCP); Central Panhandle Aquatic Preserves. [Central Panhandle Aquatic Preserves Seagrass Monitoring](#). (2023).
8. Florida Department of Environmental Protection (DEP); Office of Resilience and Coastal Protection (RCP); Central Panhandle Aquatic Preserves; Florida Fish and Wildlife Conservation Commission (FWC). [Franklin County Coastal Waters Seagrass Monitoring](#). (2017).