

# Rocky Bayou State Park Aquatic Preserve

## SEACAR Habitat Analyses

Last compiled on 08 January, 2025

### Contents

<b>Funding &amp; Acknowledgements</b>	<b>2</b>
<b>Threshold Filtering</b>	<b>2</b>
<b>Value Qualifiers</b>	<b>3</b>
<b>Water Column</b>	<b>5</b>
<b>Seasonal Kendall-Tau Analysis</b>	<b>5</b>
<b>Water Quality - Discrete</b>	<b>5</b>
Chlorophyll a, Corrected for Pheophytin - Discrete Water Quality . . . . .	6
Chlorophyll a, Uncorrected for Pheophytin - Discrete Water Quality . . . . .	7
Colored Dissolved Organic Matter - Discrete Water Quality . . . . .	9
Dissolved Oxygen - Discrete Water Quality . . . . .	11
Dissolved Oxygen Saturation - Discrete Water Quality . . . . .	13
pH - Discrete Water Quality . . . . .	15
Salinity - Discrete Water Quality . . . . .	17
Secchi Depth - Discrete Water Quality . . . . .	19
Total Nitrogen - Discrete Water Quality . . . . .	21
Total Phosphorus - Discrete Water Quality . . . . .	24
Turbidity - Discrete Water Quality . . . . .	26
Water Temperature - Discrete Water Quality . . . . .	28
<b>References</b>	<b>31</b>

## Funding & Acknowledgements

The data used in this analysis is from the Export Standardized Tables in the SEACAR Data Discovery Interface (DDI). Documents and information available through the SEACAR DDI are owned by the data provider(s) and users are expected to provide appropriate credit following accepted citation formats. Users are encouraged to access data to maximize utilization of gained knowledge, reducing redundant research and facilitating partnerships and scientific innovation.

With respect to documents and information available from SEACAR DDI, neither the State of Florida nor the Florida Department of Environmental Protection makes any warranty, expressed or implied, including the warranties of merchantability and fitness for a particular purpose arising out of the use or inability to use the data, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

This report was funded in part, through a grant agreement from the Florida Department of Environmental Protection, Florida Coastal Management Program, by a grant provided by the Office for Coastal Management under the Coastal Zone Management Act of 1972, as amended, National Oceanic and Atmospheric Administration. The views, statements, findings, conclusions and recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of the State of Florida, NOAA or any of their sub agencies.

Published: 2025-01-08



## Threshold Filtering

Threshold filters, following the guidance of Florida Department of Environmental Protection’s (*FDEP*) Division of Environmental Assessment and Restoration (*DEAR*) are used to exclude specific results values from the SEACAR Analysis. Based on the threshold filters, Quality Assurance / Quality Control (*QAQC*) Flags are inserted into the *SEACAR\_QAQCFlagCode* and *SEACAR\_QAQC\_Description* columns of the export data. The *Include* column indicates whether the *QAQC* Flag will also indicate that data are excluded from analysis. No data are excluded from the data export, but the analysis scripts can use the *Include* column to exclude data (1 to include, 0 to exclude).

Table 1: Continuous Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	-0.000001	50
Dissolved Oxygen Saturation	%	-0.000001	500
Salinity	ppt	-0.000001	70
Turbidity	NTU	-0.000001	4000
Water Temperature	Degrees C	-5.000000	45
pH	None	2.000000	14

Table 2: Discrete Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Ammonia, Un-ionized (NH3)	mg/L	-	-
Ammonium, Filtered (NH4)	mg/L	-	-
Chlorophyll a, Corrected for Pheophytin	ug/L	-	-
Chlorophyll a, Uncorrected for Pheophytin	ug/L	-	-
Colored Dissolved Organic Matter	PCU	-	-

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	-0.000001	25
Dissolved Oxygen Saturation	%	-0.000001	310
Fluorescent Dissolved Organic Matter	QSE	-	-
Light Extinction Coefficient	m <sup>-1</sup>	-	-
NO <sub>2</sub> + <sub>3</sub> , Filtered	mg/L	-	-
Nitrate (NO <sub>3</sub> )	mg/L	-	-
Nitrite (NO <sub>2</sub> )	mg/L	-	-
Nitrogen, organic	mg/L	-	-
Phosphate, Filtered (PO <sub>4</sub> )	mg/L	-	-
Salinity	ppt	-0.000001	70
Secchi Depth	m	0.000001	50
Specific Conductivity	mS/cm	0.005000	100
Total Kjeldahl Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Phosphorus	mg/L	-	-
Total Suspended Solids	mg/L	-	-
Turbidity	NTU	-	-
Water Temperature	Degrees C	3.000000	40
pH	None	2.000000	13

Table 3: Quality Assurance Flags inserted based on threshold checks listed in Table 1 and 2

<i>SEACAR QAQC Description</i>	<i>Include</i>	<i>SEACAR QAQCFlagCode</i>
Exceeds maximum threshold	0	2Q
Below minimum threshold	0	4Q
Within threshold tolerance	1	6Q
No defined thresholds for this parameter	1	7Q

## Value Qualifiers

Value qualifier codes included within the data are used to exclude certain results from the analysis. The data are retained in the data export files, but the analysis uses the *Include* column to filter the results.

### STORET and WIN value qualifier codes

Value qualifier codes from *STORET* and *WIN* data are examined with the database and used to populate the *Include* column in data exports.

Table 4: Value Qualifier codes excluded from analysis

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>MDL</i>	<i>Description</i>
STORET-WIN	H	0	0	Value based on field kit determination; results may not be accurate
STORET-WIN	J	0	0	Estimated value
STORET-WIN	V	0	0	Analyte was detected at or above method detection limit
STORET-WIN	Y	0	0	Lab analysis from an improperly preserved sample; data may be inaccurate

### Discrete Water Quality Value Qualifiers

The following value qualifiers are highlighted in the Discrete Water Quality section of this report. An exception is made for **Program 476 - Charlotte Harbor Estuaries Volunteer Water Quality Monitoring Network** and data flagged with Value Qualifier **H** are included for this program only.

**H** - Value based on field kit determination; results may not be accurate. This code shall be used if a field screening test (e.g., field gas chromatograph data, immunoassay, or vendor-supplied field kit) was used to generate the value and the field kit or method has not been recognized by the Department as equivalent to laboratory methods.

**I** - The reported value is greater than or equal to the laboratory method detection limit but less than the laboratory practical quantitation limit.

**Q** - Sample held beyond the accepted holding time. This code shall be used if the value is derived from a sample that was prepared or analyzed after the approved holding time restrictions for sample preparation or analysis.

**S** - Secchi disk visible to bottom of waterbody. The value reported is the depth of the waterbody at the location of the Secchi disk measurement.

**U** - Indicates that the compound was analyzed for but not detected. This symbol shall be used to indicate that the specified component was not detected. The value associated with the qualifier shall be the laboratory method detection limit. Unless requested by the client, less than the method detection limit values shall not be reported

**Systemwide Monitoring Program (SWMP) value qualifier codes**

Value qualifier codes from the *SWMP* continuous program are examined with the database and used to populate the *Include* column in data exports. *SWMP* Qualifier Codes are indicated by *QualifierSource=SWMP*.

Table 5: SWMP Value Qualifier codes

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>Description</i>
SWMP	-1	Yes	Optional parameter not collected
SWMP	-2	No	Missing data
SWMP	-3	No	Data rejected due to QA/QC
SWMP	-4	No	Outside low sensor range
SWMP	-5	No	Outside high sensor range
SWMP	0	Yes	Passed initial QA/QC checks
SWMP	1	No	Suspect data
SWMP	2	Yes	Reserved for future use
SWMP	3	Yes	Calculated data: non-vented depth/level sensor correction for changes in barometric pressure
SWMP	4	Yes	Historical: Pre-auto QA/QC
SWMP	5	Yes	Corrected data

## Water Column

The water column habitat extends from the water's surface to the bottom sediments, and it's where fish, dolphins, crabs and people swim! So much life makes its home in the water column that the health of marine and coastal ecosystems, as well as human economies, depend on the condition of this vulnerable habitat. Local patterns of rainfall, temperature, winds and currents can rapidly change the condition of the water column, while global influences such as [El Niño/La Niña](#), large-scale fluctuation in sea temperatures and climate change can have long-term effects. Inputs from the prosperity of our day-to-day lives including farming, mining and forestry, and emissions from power generation, automobiles and water treatment can also alter the health of the water column. Acting alone or together, each input can have complex and lasting effects on habitats and ecosystems.

SEACAR evaluates water column health with several essential parameters. These include nutrient surveys of nitrogen and phosphorus, and water quality assessments of salinity, dissolved oxygen, pH, and water temperature. Water clarity is evaluated with Secchi depth, turbidity, levels of chlorophyll a, total suspended solids, and colored dissolved organic matter. Additionally, the richness of nekton is indicated by the abundance of free-swimming fishes and macroinvertebrates like crabs and shrimps.

## Seasonal Kendall-Tau Analysis

Indicators must have a minimum of five to ten years, depending on the habitat, of data within the geographic range of the analysis to be included in the analysis. Ten years of data are required for discrete parameters, and five years of data are required for continuous parameters. If there are insufficient years of data, the number of years of data available will be noted and labeled as “insufficient data to conduct analysis”. Further, for the preferred Seasonal Kendall-Tau test, there must be data from at least two months in common across at least two consecutive years within the RCP managed area being analyzed. Values that pass both of these tests will be included in the analysis and be labeled as *Use\_In\_Analysis* = **TRUE**. Any that fail either test will be excluded from the analyses and labeled as *Use\_In\_Analysis* = **FALSE**. The points for all Water Column plots displayed in this section are monthly averages. Trend significance will be denoted as “Significant Trend” (when  $p < 0.05$ ), or “Non-significant Trend” (when  $p \geq 0.05$ ). Any parameters with insufficient data to perform Seasonal Kendall-Tau test will have their monthly averages plotted without a corresponding trend line.

## Water Quality - Discrete

The following files were used in the discrete analysis:

- *Combined\_WQ\_WC\_NUT\_Chlorophyll\_a\_corrected\_for\_pheophytin-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Chlorophyll\_a\_uncorrected\_for\_pheophytin-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Colored\_dissolved\_organic\_matter\_CDOM-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Dissolved\_Oxygen-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Dissolved\_Oxygen\_Saturation-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_pH-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Salinity-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Secchi\_Depth-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Total\_Nitrogen-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Total\_Phosphorus-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Total\_Suspended\_Solids\_TSS-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Turbidity-2024-Dec-08.txt*
- *Combined\_WQ\_WC\_NUT\_Water\_Temperature-2024-Dec-08.txt*

# Chlorophyll a, Corrected for Pheophytin - Discrete Water Quality

## Seasonal Kendall-Tau Trend Analysis




Figure 1: Seasonal Kendall-Tau Results for Chlorophyll a, Corrected for Pheophytin - Discrete

Table 6: Seasonal Kendall-Tau Trend Analysis for Chlorophyll a, Corrected for Pheophytin

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	103	13	3	TRUE	-0.1845	0.0504	-0.025	3.2357	7.7485	0.7356	0




Figure 2: Map showing location of Discrete sampling sites for Chlorophyll a, Corrected for Pheophytin. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 7: Programs contributing data for Chlorophyll a, Corrected for Pheophytin

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	60	2012	2017
537	43	2018	2024

**Program names:**

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>  
 5002 - Florida STORET / WIN<sup>2</sup>

**Chlorophyll a, Uncorrected for Pheophytin - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**

Chlorophyll a, Uncorrected for Pheophytin - Discrete, All Depths  
 Rocky Bayou State Park Aquatic Preserve




Figure 3: Seasonal Kendall-Tau Results for Chlorophyll a, Uncorrected for Pheophytin - Discrete

Table 8: Seasonal Kendall-Tau Trend Analysis for Chlorophyll a, Uncorrected for Pheophytin

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	452	24	3.5	TRUE	-0.017	0.78	0	3.4974	10.7687	0.4628	0






Figure 4: Map showing location of Discrete sampling sites for Chlorophyll a, Uncorrected for Pheophytin. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 9: Programs contributing data for Chlorophyll a, Uncorrected for Pheophytin

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
514	204	2001	2019
5002	183	2001	2017
537	52	2018	2024
103	13	2020	2021

**Program names:**

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>3</sup>

514 - Florida LAKEWATCH Program<sup>4</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

**Colored Dissolved Organic Matter - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 5: Seasonal Kendall-Tau Results for Colored Dissolved Organic Matter - Discrete

Table 10: Seasonal Kendall-Tau Trend Analysis for Colored Dissolved Organic Matter

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	128	24	13.5	TRUE	0.0916	0.3155	0.1818	9.6216	9.3936	0.5856	0




Figure 6: Map showing location of Discrete sampling sites for Colored Dissolved Organic Matter. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 11: Programs contributing data for Colored Dissolved Organic Matter

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
514	108	2001	2019
537	20	2019	2024

**Program names:**

514 - Florida LAKEWATCH Program<sup>4</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

**Dissolved Oxygen - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 7: Seasonal Kendall-Tau Results for Dissolved Oxygen - Discrete

Table 12: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	770	26	6.61	TRUE	0.1158	0.0082	0.0224	5.5927	12.511	0.3265	1




Figure 8: Map showing location of Discrete sampling sites for Dissolved Oxygen. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 13: Programs contributing data for Dissolved Oxygen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	387	2008	2024
5002	347	2001	2017
69	20	1994	2022
103	12	2020	2021
102	4	1994	1994

**Program names:**

- 69 - Fisheries-Independent Monitoring (FIM) Program<sup>5</sup>
- 102 - National Status and Trends Mussel Watch<sup>6</sup>
- 103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>3</sup>
- 537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>
- 5002 - Florida STORET / WIN<sup>2</sup>

**Dissolved Oxygen Saturation - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 9: Seasonal Kendall-Tau Results for Dissolved Oxygen Saturation - Discrete

Table 14: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen Saturation

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	731	25	89.3	TRUE	0.1513	0.0009	0.4625	69.8352	9.6384	0.5632	1




Figure 10: Map showing location of Discrete sampling sites for Dissolved Oxygen Saturation. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 15: Programs contributing data for Dissolved Oxygen Saturation

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	383	2008	2024
5002	348	2000	2017

**Program names:**

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>  
 5002 - Florida STORET / WIN<sup>2</sup>

**pH - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 11: Seasonal Kendall-Tau Results for pH - Discrete

Table 16: Seasonal Kendall-Tau Trend Analysis for pH

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	766	27	7.9	TRUE	0.049	0.2436	0.003	7.841	9.9314	0.5366	0






Figure 12: Map showing location of Discrete sampling sites for pH. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 17: Programs contributing data for pH

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	387	2008	2024
5002	347	2000	2017
69	20	1994	2022
103	12	2020	2021

**Program names:**

69 - Fisheries-Independent Monitoring (FIM) Program<sup>5</sup>

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>3</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

**Salinity - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 13: Seasonal Kendall-Tau Results for Salinity - Discrete

Table 18: Seasonal Kendall-Tau Trend Analysis for Salinity

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	756	27	22	TRUE	0.0502	0.2844	0.0397	21.2907	8.9419	0.6273	0




Figure 14: Map showing location of Discrete sampling sites for Salinity. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 19: Programs contributing data for Salinity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	384	2008	2024
5002	348	2000	2017
69	20	1994	2022
102	4	1994	1994

**Program names:**

69 - Fisheries-Independent Monitoring (FIM) Program<sup>5</sup>

102 - National Status and Trends Mussel Watch<sup>6</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

**Secchi Depth - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 15: Seasonal Kendall-Tau Results for Secchi Depth - Discrete

Table 20: Seasonal Kendall-Tau Trend Analysis for Secchi Depth

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
Surface	458	25	2.4384	TRUE	-0.1678	0.0003	-0.0254	2.7703	9.0299	0.6191	-1




Figure 16: Map showing location of Discrete sampling sites for Secchi Depth. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 21: Programs contributing data for Secchi Depth

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	229	2001	2024
514	201	2001	2019
5002	16	2013	2017
69	12	1995	2022

**Program names:**

69 - Fisheries-Independent Monitoring (FIM) Program<sup>5</sup>

514 - Florida LAKEWATCH Program<sup>4</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

**Total Nitrogen - Discrete Water Quality**

**Total Nitrogen Calculation:**

The logic for calculated Total Nitrogen was provided by Kevin O'Donnell and colleagues at FDEP (with the help of Jay Silvanima, Watershed Monitoring Section). The following logic is used, in this order, based on the availability of specific nitrogen components.

- 1)  $TN = TKN + NO3O2$ ;
- 2)  $TN = TKN + NO3 + NO2$ ;
- 3)  $TN = ORGN + NH4 + NO3O2$ ;
- 4)  $TN = ORGN + NH4 + NO2 + NO3$ ;
- 5)  $TN = TKN + NO3$ ;
- 6)  $TN = ORGN + NH4 + NO3$ ;

Additional Information:

- Rules for use of sample fraction:
  - Florida Department of Environmental Protection (FDEP) report that if both “Total” and “Dissolved” components are reported, only “Total” is used. If the total is not reported, then the dissolved components are used as a best available replacement.
  - Total nitrogen calculations are done using nitrogen components with the same sample fraction, nitrogen components with mixed total/dissolved sample fractions are not used. In other words, total nitrogen can be calculated when TKN and NO3O2 are both total sample fractions, or when both are dissolved sample fractions. *Future calculations of total nitrogen values may be based on components with mixed sample fractions.*
- Values inserted into data:
  - ParameterName = “Total Nitrogen”
  - SEACAR\_QAQCFlagCode = “1Q”
  - SEACAR\_QAQC\_Description = “SEACAR Calculated”

### Seasonal Kendall-Tau Trend Analysis




Figure 17: Seasonal Kendall-Tau Results for Total Nitrogen - Discrete

Table 22: Seasonal Kendall-Tau Trend Analysis for Total Nitrogen

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	635	24	0.2	TRUE	0.1611	0.0002	0.0022	0.1462	6.4986	0.8381	1




Figure 18: Map showing location of Discrete sampling sites for Total Nitrogen. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 23: Programs contributing data for Total Nitrogen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	246	2001	2024
514	203	2001	2019
5002	186	2001	2017

**Program names:**

514 - Florida LAKEWATCH Program<sup>4</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

**Total Phosphorus - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**



Total Phosphorus - Discrete, All Depths  
Rocky Bayou State Park Aquatic Preserve




Figure 19: Seasonal Kendall-Tau Results for Total Phosphorus - Discrete

Table 24: Seasonal Kendall-Tau Trend Analysis for Total Phosphorus

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	465	24	0.007	TRUE	0.2622	0	0.0002	0.0035	6.9238	0.8052	1




Figure 20: Map showing location of Discrete sampling sites for Total Phosphorus. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 25: Programs contributing data for Total Phosphorus

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	246	2001	2024
514	203	2001	2019
5002	17	2013	2017
103	8	2020	2021

**Program names:**

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>3</sup>

514 - Florida LAKEWATCH Program<sup>4</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

**Turbidity - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**




Figure 21: Seasonal Kendall-Tau Results for Turbidity - Discrete

Table 26: Seasonal Kendall-Tau Trend Analysis for Turbidity

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	222	11	1	TRUE	-0.2377	0.0034	-0.2134	3.4417	15.4432	0.1631	-2




Figure 22: Map showing location of Discrete sampling sites for Turbidity. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 27: Programs contributing data for Turbidity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	375	2008	2024
5002	304	2001	2017
103	12	2020	2021

**Program names:**

- 103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX<sup>3</sup>
- 537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>
- 5002 - Florida STORET / WIN<sup>2</sup>

**Water Temperature - Discrete Water Quality**

**Seasonal Kendall-Tau Trend Analysis**

Water Temperature - Discrete, All Depths  
 Rocky Bayou State Park Aquatic Preserve




Figure 23: Seasonal Kendall-Tau Results for Water Temperature - Discrete

Table 28: Seasonal Kendall-Tau Trend Analysis for Water Temperature

RelativeDepth	N-Data	N-Years	Median	Independent	tau	p	SennSlope	SennIntercept	ChiSquared	pChiSquared	Trend
All	576	19	24.3333	TRUE	0.1517	0.0063	0.0754	22.499	12.9782	0.2948	1




Figure 24: Map showing location of Discrete sampling sites for Water Temperature. The bubble size on the maps below reflect the amount of data available at each sampling site.

Table 29: Programs contributing data for Water Temperature

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
537	501	2008	2024
5002	39	2013	2017
69	20	1994	2022
103	12	2020	2021
102	4	1994	1994

**Program names:**

69 - Fisheries-Independent Monitoring (FIM) Program<sup>5</sup>

102 - National Status and Trends Mussel Watch<sup>6</sup>

103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX<sup>3</sup>

537 - Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring<sup>1</sup>

5002 - Florida STORET / WIN<sup>2</sup>

## References

1. Choctawhatchee Basin Alliance. [Choctawhatchee Basin Alliance Living Shorelines Oyster Reef Monitoring](#). (2024).
2. Florida Department of Environmental Protection (DEP). [Florida STORET / WIN](#). (2024).
3. U.S. Environmental Protection Agency (EPA). [EPA STOrage and RETrieval Data Warehouse \(STORET\)/WQX](#). (2023).
4. University of Florida (UF); Institute of Food and Agricultural Sciences. [Florida LAKEWATCH Program](#). (2024).
5. Florida Fish and Wildlife Conservation Commission (FWC). [Fisheries-Independent Monitoring \(FIM\) Program](#). (2022).
6. National Oceanic and Atmospheric Administration (NOAA); Center for Coastal Monitoring and Assessment. [National Status and Trends Mussel Watch](#). (2000).